



**Defense Threat Reduction Agency
8725 John J. Kingman Road, MS-6201
Fort Belvoir, VA 22060-6201**



DTRA-TR-17-026 (R1)

TECHNICAL REPORT

Publicly Released Prompt Radiation Spectra Suitable for Nuclear Detonation Simulations, Revision 1

DISTRIBUTION A. Approved for public release; distribution is unlimited.



December 2017

Prepared by:

Applied Research Associates, Inc.
801 N. Quincy Street
Suite 700
Arlington, VA 22203

This page intentionally left blank.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503.</small> PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 12-01-2017		2. REPORT TYPE Technical Report		3. DATES COVERED (From – To)	
4. TITLE AND SUBTITLE Publicly Released Prompt Radiation Spectra Suitable for Nuclear Detonation Simulations, Revision 1				5a. CONTRACT NUMBER HDTRA1-14-D-0003; 0005	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Kevin Kramer, Tyler Dant, Andy Li, and Kyle Millage				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Applied Research Associates, Inc. 801 N. Quincy St., Suite 700 Arlington, VA 22203				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Defense Threat Reduction Agency 8725 John J. Kingman Road, MS 6201 Fort Belvoir, VA 22060-6201				10. SPONSOR ACRONYM(S) DTRA RD	
				11. SPONSOR'S REPORT NUMBER(S) DTRA-TR-17-026 (R1)	
12. DISTRIBUTION / AVAILABILITY STATEMENT DISTRIBUTION A: Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT This technical report describes unclassified source leakage spectra that can be used to simulate a nuclear device or weapon in a radiation transport model. The subsequent radiation transport model results are essential input for civilian emergency planning, consequence assessment, and medical countermeasure development. This report suggests that the four sets of neutron and photon spectra are the most suitable for simulations: the two spectra sets from Hiroshima and Nagasaki fission weapons and two spectra sets from DNA 4267F. These were chosen by the authors due to the relative completeness of their quantitative description. An associated spreadsheet document is available with this technical report that provides numerical data sets to support radiation transport modeling.					
15. SUBJECT TERMS Radiation Transport Nuclear Detonation Initial Radiation Urban Terrain Dose Estimate					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Unlimited	18. NUMBER OF PAGES 21	19a. NAME OF RESPONSIBLE PERSON Paul K. Blake, PhD
a. REPORT U	b. ABSTRACT U	a. THIS PAGE U			19b. PHONE NUMBER (include area code) 703 767-3433

UNIT CONVERSION TABLE			
U.S. customary units to and from international units of measurement*			
U.S. Customary Units	Multiply by		International Units
	 Divide by†		
Length/Area/Volume			
inch (in)	2.54	× 10 ⁻²	meter (m)
foot (ft)	3.048	× 10 ⁻¹	meter (m)
yard (yd)	9.144	× 10 ⁻¹	meter (m)
mile (mi, international)	1.609 344	× 10 ³	meter (m)
mile (nmi, nautical, U.S.)	1.852	× 10 ³	meter (m)
barn (b)	1	× 10 ⁻²⁸	square meter (m ²)
gallon (gal, U.S. liquid)	3.785 412	× 10 ⁻³	cubic meter (m ³)
cubic foot (ft ³)	2.831 685	× 10 ⁻²	cubic meter (m ³)
Mass/Density			
pound (lb)	4.535 924	× 10 ⁻¹	kilogram (kg)
atomic mass unit (AMU)	1.660 539	× 10 ⁻²⁷	kilogram (kg)
pound-mass per cubic foot (lb ft ⁻³)	1.601 846	× 10 ¹	kilogram per cubic meter (kg m ⁻³)
pound-force (lbf avoirdupois)	4.448 222		newton (N)
Energy/Work/Power			
electronvolt (eV)	1.602 177	× 10 ⁻¹⁹	joule (J)
Erg	1	× 10 ⁻⁷	joule (J)
kiloton (kT) (TNT equivalent)	4.184	× 10 ¹²	joule (J)
British thermal unit (Btu) (thermochemical)	1.054 350	× 10 ³	joule (J)
foot-pound-force (ft lbf)	1.355 818		joule (J)
calorie (cal) (thermochemical)	4.184		joule (J)
Pressure			
atmosphere (atm)	1.013 250	× 10 ⁵	pascal (Pa)
pound force per square inch (psi)	6.984 757	× 10 ³	pascal (Pa)
Temperature			
degree Fahrenheit (°F)	[T(°F) – 32]/1.8		degree Celsius (°C)
degree Fahrenheit (°F)	[T(°F) + 459.67]/1.8		kelvin (K)
Radiation			
curie (Ci) (activity of radionuclides)	3.7	× 10 ¹⁰	s ^{-1‡}
air exposure (roentgen)	2.579 760	× 10 ⁻⁴	coulomb per kilogram (C kg ⁻¹)
absorbed dose (rad)	1	× 10 ⁻²	J kg ^{-1§}
equivalent and effective dose (rem)	1	× 10 ⁻²	J kg ^{-1**}

* Specific details regarding the implementation of SI units may be viewed at <http://www.bipm.org/en/si/>.

† Multiply the U.S. customary unit by the factor to get the international unit. Divide the international unit by the factor to get the U.S. customary unit.

‡ The special name for the SI unit of the activity of a radionuclide is the becquerel (Bq). (1 Bq = 1 s⁻¹).

§ The special name for the SI unit of absorbed dose is the gray (Gy). (1 Gy = 1 J kg⁻¹).

** The special name for the SI unit of equivalent and effective dose is the sievert (Sv). (1 Sv = 1 J kg⁻¹).

Table of Contents

LIST OF FIGURES.....	II
LIST OF TABLES	III
REVISION NOTES	IV
ACKNOWLEDGEMENTS	V
EXECUTIVE SUMMARY	1
1 INTRODUCTION	2
2 DESCRIPTION OF PROMPT RADIATION SPECTRA	3
3 DISCUSSION OF SPECTRA	5
4 CONCLUSIONS ON USE OF SPECTRA	7
BIBLIOGRAPHY	11
ABBREVIATIONS, ACRONYMS, AND SYMBOLS.....	12

List of Figures

Figure 1. Neutron spectra from fission sources	7
Figure 2. Fission neutron spectra with values greater than 0.1 MeV	8
Figure 3. Neutron spectra from thermonuclear sources	8
Figure 4. Thermonuclear neutron spectra with energies greater than 0.1 MeV	9
Figure 5. Photon spectra from fission sources.....	9
Figure 6. Photon spectra from thermonuclear sources.....	10

List of Tables

Table 1 - List of unclassified spectra	3
Table 2. Total number of particles used for plotting	6
Table 3. Suggested spectra for simulation of leakage radiation for a nuclear device	7

Revision Notes

The first purpose of this revision is to change the primary reference for the 'Little Boy' and 'Fat Man' spectra from LA-UR-01-6594, a publication from Los Alamos National Laboratories by White, Whalen and Heath (White, Whalen, & Heath, 2001), to the more recent publication LA-UR-13-26113 (Holmes & White, 2013). The gamma yield for the 'Fat Man' spectra is different in LA-UR-01-6594 and is inconsistent with the Hiroshima and Nagasaki Dosimetry (Young, R.W. & Kerr, G.W. [Eds.], 2005). While this report is focused on the spectra and not the gamma yield, the authors would prefer the primary reference be consistent with the official Hiroshima and Nagasaki dosimetry. The accompanying spreadsheet has also been updated to reflect this change in emphasis.

The second purpose of this revision is to include a description of the White Sands Missile Range Fast Burst Reactor from a publication by Flanders and Sparks (Flanders & Sparks, 1989) to our list of publically available fission sources.

Acknowledgements

We would like to thank Dr. Blake of the Defense Threat Reduction Agency (DTRA) for sponsoring and encouraging this work. We would also like to thank Amy Curcio for help with editing and formatting. We would like to thank Dr's Stephen Egbert, David Schauer, and Lisa Andivahis for providing comments and suggesting useful references.

Executive Summary

This technical report contains a set of neutron and photon spectra that can be used for simulating the prompt radiation from a nuclear device or weapon in a radiation transport model. Applied Research Associates, Inc. (ARA) collected these weapon outputs to improve simulations of the prompt radiation of nuclear devices through complex urban environments. The simulations are for emergency planning, consequence assessment, and medical countermeasure development by the civilian community.

After listing twelve different sets of data, this technical report suggests that four sets of neutron and photon spectra are the most suitable for simulations: the two sets of spectra from Hiroshima and Nagasaki fission weapons and two sets of spectra from DNA 4267F. These were chosen by the authors due to the relative completeness of their quantitative description.

1 Introduction

As part of its mission to safeguard against weapons of mass destruction (WMD), the Defense Threat Reduction Agency (DTRA) supports the development of capabilities to reduce, eliminate and counter WMD threats and mitigate their effects. Applied Research Associates (ARA) was tasked by DTRA to support this mission by providing publicly available nuclear detonation radiation spectra essential for the civilian community in emergency planning, consequence assessment, and medical countermeasure development. This technical report describes different unclassified leakage spectra that can be used to simulate a nuclear device or weapon in a radiation transport model.

ARA has developed capabilities to simulate the prompt radiation of nuclear devices in a three-dimensional Monte Carlo radiation transport model. These models include 3-D representation of urban geometries complete with internal structure and external walls. ARA uses the models to improve consequence assessment analysis to help inform nuclear terrorism response planning communication to the public but respects the sensitive nature of nuclear weapon effects simulations. This technical report describes various prompt radiation spectra and attempts to sort out the ones that can be both meaningful representations of possible devices and are also available for public release.

An associated spreadsheet is available with this technical report that provides numerical data sets to support radiation transport modeling.

2 Description of Prompt Radiation Spectra

The prompt radiation spectrum is the neutron and photon radiation released in the short time, much less than a microsecond, after detonation (Glasstone, 1977). The detonation releases both neutron and photons in this time; however, the neutron emission dominates the photon emission. During the Hiroshima and Nagasaki bombings, the prompt radiation contributed from 40% to 70% of the free-in-air dose depending on distance from epicenter (Young, R.W. & Kerr, G.W. [Eds.], 2005). Some of the leakage spectra for the photon spectrum listed are not available or contain more sensitive information than the neutron spectrum. Table 1 lists various spectra that are available. Note, all the spectra listed are unclassified.

Table 1 - List of unclassified spectra

<i>Source</i>	<i>Description</i>	<i>Both Neutron and Photon Spectra?</i>
Holmes & White 2013 (Little Boy)	Analysis of ^{235}U Hiroshima fission weapon	Yes
Holmes & White 2013 (Fat Man)	Analysis of ^{239}Pu Nagasaki fission weapon	Yes
Whalen 1983 (Little Boy)	Analysis of ^{235}U Hiroshima fission weapon	Yes
Whalen 1983 (Fat Man)	Analysis of ^{239}Pu Nagasaki fission weapon	Yes
Glasstone & Dolan 1977 (Fission)	Neutron fission spectrum from "The Effects of Nuclear Weapons"	No
Glasstone & Dolan 1977 (Thermonuclear)	Neutron spectrum from thermonuclear device from "The Effects of Nuclear Weapons"	No
Terrell 1990	MCNP calculation of idealized Godiva device	Yes
Kaul & Egbert 2000	Neutron spectrum from unshielded Army Pulse Reactor Division reactor	Yes
Auxier, et al. 1972 (Low, Intermediate, High)	Suggested neutron spectra for high-, intermediate- and low-yield thermonuclear weapons for initial radiation shielding calculations	No
Gritzner, et al. 1976 (EM-1, Low, Henre)	Source spectra from potential tactical nuclear weapons and the Henre experiment	Yes

Barnes, et al. 1991	Estimated spectrum for a boosted fission weapon	No
Flanders & Sparks 1987	Source spectra for White Sands Missile Range Fast Burst Reactor	No

3 Discussion of Spectra

There are four categories of neutron spectra presented: “Little Boy”, “Fat Man”, unshielded or bare sphere, and thermonuclear. “Little Boy” is a gun-assembled weapon while “Fat Man” is a high-explosive assembled weapon, but since there aren’t other similar spectra in the public domain, the authors will refer to them by their names. The Radiation Effects Research Foundation (RERF) has extensively modeled the “Little Boy” and “Fat Man” weapon spectra from the Hiroshima and Nagasaki nuclear detonations, respectively, and the radiation spectra are fairly detailed (Young, R.W. & Kerr, G.W. [Eds.], 2005). The differences between the two spectra show the sensitivity of these devices to their design. The spectra for “Little Boy” are known to have a significant angular dependence; however, for the applications of emergency planning, consequence assessment, and medical countermeasure development isotropic sources are more suitable. The isotropic, angular-average version of the spectrum is plotted in this report and provided in the accompanying spreadsheet. A 2013 LANL report (Holmes & White, 2013) presents a new look at the “Little Boy” and “Fat Man” outputs, but the published output spectra did not change from the 2005 DS02 report.

The Glasstone & Dolan 1977 fission spectra is a calculation of a bare sphere undergoing ^{235}U fission. Glasstone & Dolan 1977 includes a thermonuclear spectrum as well. The APRD (the Army Pulse Radiation Division) report (Kaul & Egbert, 2000) contains a calculation based on experimental data from an unshielded reactor. Flanders and Sparks have a similar calculation for the White Sands Missile Range Fast Burst Reactor (Flanders and Sparks, 1989). The Terrell report details an MCNP (Monte Carlo N-Particle radiation transport code) calculation of an idealized Godiva device with 93.77% ^{235}U . The Oak Ridge report (Auxier, et al., 1976) includes three “typical” thermonuclear neutron spectra created from transport calculations and suggests the intermediate one (300 kT) as the standard. A second Oak Ridge report (Barnes, et al. 1991) contains an estimated neutron spectrum for a boosted fission device. The Defense Nuclear Agency report 4267F (Gritzner, et al., 1972) contains three source spectra: an unshielded fission source, a thermonuclear spectrum from a hypothetical low-yield weapon and a spectrum from the reactor employed in the Henre experiment.

All the fission neutron spectra identified in Table 1 are plotted in Figure 1. Figure 2 is the same plot, but showing only the neutron energies above 0.1 MeV. Figure 3 shows the thermonuclear neutron spectra with Figure 4 showing the same, but with the domain again limited to neutron energies above 0.1 MeV. Figure 5 shows the available primary photon spectra for the corresponding fission sources. Figure 6 shows the primary photon spectra for the two thermonuclear spectra in DNA 4267F.

Each data set has its own method of binning particle energy; therefore, for purposes of comparison, each histogram bin is divided by the difference in energy between bin divisions. The authors used total neutrons per kT for normalization. However, data sets are frequently published as unnormalized fractions. In those cases, the authors of this report used neutrons (or photons) per kT from one of the Glasstone spectra or from one of the DNA 4267F spectra. The normalizations are listed in Table 2.

The authors extracted the total number of neutrons per kT for both of the Glasstone & Dolan 1977 spectra by digitizing the plots in Figure 8.116a and Figure 8.116b of Glasstone & Dolan 1977 and summing the values of each histogram bin. This quantity is only used for comparison.

The details of each spectrum are contained in a spreadsheet which is available on request.

Table 2. Total number of particles used for plotting

Source Spectra	Particle Type	Source	Total Particles per kT for comparison normalization
Little Boy	Neutron	Holmes & White 2013	1.065×10^{23}
Fat Man	Neutron	Holmes & White 2013	1.590×10^{23}
APRD reactor	Neutron	Glasstone & Dolan 1977	7.760×10^{22}
WSMR reactor	Neutron	Glasstone & Dolan 1977	7.760×10^{22}
Glasstone (Fission)	Neutron	Glasstone & Dolan 1977	7.760×10^{22}
Terrell	Neutron	Glasstone & Dolan 1977	7.760×10^{22}
DNA 4267F (EM-1 Fission)	Neutron	Gritzner, et al. 1976	1.00×10^{23}
Glasstone (Thermonuclear)	Neutron	Glasstone & Dolan 1977	1.445×10^{23}
ORNL-TM-11775 (Boosted Fission)	Neutron	Barnes, et al. 1991	3.102×10^{23}
ORNL-TM-3396 (High Yield Thermonuclear)	Neutron	Glasstone & Dolan 1977	1.445×10^{23}
ORNL-TM-3396 (Intermediate Yield Thermonuclear)	Neutron	Glasstone & Dolan 1977	1.445×10^{23}
ORNL-TM-3396 (Low Yield Thermonuclear)	Neutron	Glasstone & Dolan 1977	1.445×10^{23}
DNA 4267F (Low Yield Thermonuclear)	Neutron	Gritzner, et al. 1976	2.00×10^{23}
DNA 4276F (Henre)	Neutron	Gritzner, et al. 1976	1.00×10^{24}
Little Boy	Photon	Holmes & White 2013	4.014×10^{21}
Fat Man	Photon	Holmes & White 2013	5.43×10^{22}
Terrell	Photon	Gritzner, et al. 1976	3.00×10^{22}
APRD reactor	Photon	Gritzner, et al. 1976	3.00×10^{22}
DNA 4276F (EM-1 Fission)	Photon	Gritzner, et al. 1976	3.00×10^{22}
DNA 4276F (Low Yield Thermonuclear)	Photon	Gritzner, et al. 1976	7.0×10^{22}
DNA 4276F (Henre)	Photon	Gritzner, et al. 1976	1.0×10^{23}

4 Conclusions on Use of Spectra

The suggested spectra for use in simulations of prompt radiation from nuclear devices intended for public release are shown in Table 3. The authors chose these spectra because of the resolution and range, possessing information about source particles per kT normalizations and having a corresponding primary photon spectrum.

Table 3. Suggested spectra for simulation of leakage radiation for a nuclear device

Type of Spectrum	Neutron Source	Photon Source
“Little Boy”-type spectrum	Holmes & White 2013	Holmes & White 2013
“Fat Man”-type spectrum	Holmes & White 2013	Holmes & White 2013
Unshielded device spectrum	Gritzner, et al. 1976 (EM-1 Fission)	Gritzner, et al. 1976 (EM-1 Fission)
Thermonuclear spectrum	Gritzner, et al. 1976 (Low Yield Thermonuclear)	Gritzner, et al. 1976 (Low Yield Thermonuclear)




Figure 1. Neutron spectra from fission sources




Figure 2. Fission neutron spectra with values greater than 0.1 MeV




Figure 3. Neutron spectra from thermonuclear sources




Figure 4. Thermonuclear neutron spectra with energies greater than 0.1 MeV




Figure 5. Photon spectra from fission sources




Figure 6. Photon spectra from thermonuclear sources

Bibliography

- Auxier, J. A., Burson, Z. G., French, R. L., Haywood, F. F., Mooney, L. G., & Stroker, E. A. (1972). *Nuclear Weapons Free-Field Environment Recommended for Initial Radiation Shielding Calculations*. ORNL-TM-3396, Oak Ridge, TN: Oak Ridge National Laboratories.
- Barnes, J. M., Johnson, J. O., Burns, T. J., and Drischler, J. D. (1991). *Nuclear Vulnerability of the U.S. M60A 1 Tank in an Initial Radiation Environment: Mash Code System Analysis*. ORNL-TM-11775, Oak Ridge, TN: Oak Ridge National Laboratories.
- Flanders, T. M. and Sparks, M. H. (1989). *Monte Carlo Calculations of the Neutron Environment Produced by the White Sands Missile Range Fast Burst Reactor*. Nuclear Science and Engineering. Vol. 103, Num. 3, pp. 265-275.
- Glasstone, S. & Dolan, P.J. (1977). *The Effects of Nuclear Weapons, 3rd Edition*. United States Department of Defense and United States Department of Energy.
- Gritzner, M. L., Albert, T. E., Straker, E. A., & Smith, H. T. (1976). *Radiation Environments from Tactical Nuclear Weapons*. DNA 4267F, Washington, DC: Defense Nuclear Agency.
- Holmes, R. L. and White, Stephen W. (2013). *Output of Fat Man and Little Boy Devices*. LA-UR-13-26113. Los Alamos, NM: Los Alamos National Laboratory. (Also, published as DTRA-IR-13-67)
- Kaul, D. and Egbert, S. (2000). *Radiation Environment Program (REP) 1992 to 1994*. DNA001-92-C-0082, Dulles, VA: Defense Threat Reduction Agency.
- Terrell, J. (1990). *Gamma-Ray and Neutron Leakage Spectra Calculated for Unshielded Reactors*. LA-11807, Los Alamos, NM: Los Alamos National Laboratory.
- Whalen, P. (1983). *Source Terms for the Initial Radiation*. LA-UR-83-198. US-Japan Joint Workshop on the Reassessment of the A-Bomb Radiation Dosimetry in Hiroshima and Nagasaki, Los Alamos, NM, Los Alamos National Laboratory.
- White, S., Whalen, P. and Heath, Alexandra, R. (2001) *Source Calculations for the US-Japan Dosimetry Working Group*. LA-UR-01-6594, Los Alamos National Laboratory, Los Alamos, NM: Los Alamos National Laboratory.
- Young, R.W. & Kerr, G.W. (Eds.) (2005). *Reassessment of the Atomic Bomb Radiation Dosimetry for Hiroshima and Nagasaki—Dosimetry System 2002*. Hiroshima, Japan: Radiation Effects Research Foundation.

Abbreviations, Acronyms, and Symbols

ARA	Applied Research Associates, Inc.
APRD	Army Pulse Reactor Division
ATR6	Air-Transport of Radiation Version 6
DNA	Defense Nuclear Agency
DS02	Hiroshima and Nagasaki Dosimetry System 2002
DTRA	Defense Threat Reduction Agency (United States)
HSRD	Human Survivability Research and Development Integrated Program Team
EM-1	Effects Manual volume 1
FOUO	For Official Use Only
IPT	Integrated Program Team
kT	Kiloton
MCNP	Monte Carlo N-Particle radiation transport software
mole	A number of constituent particles equal to the Avogadro constant (6.022×10^{23})
ORNL	Oak Ridge National Laboratories
RERF	Radiation Effects Research Foundation
TR	Technical Report
U.S.	United States